

Improving Survivorship for Cancer Patients


Latest Research Shows Lymphedema is a Risk of Cancer Treatment



■ 58% of cancer patients are at risk for limb lymphedema¹⁻³



- Breast cancer
- Melanoma
- Pelvic area cancer



- Lymph node surgery
- Nodal irradiation
- Taxane-based chemotherapy

Early Detection & Intervention are Key to Preventing Lymphedema



of patients did not progress to chronic lymphedema with early detection using L-Dex® and intervention through 3 years⁵

The opportunity to prevent lymphedema is before noticeable swelling exists

Testing for Early Lymphedema is Easy

L-Dex® testing with the SOZO® Digital Health Platform



Learn more
www.impedimed.com/improving-survivorship

↓
**Fast
Non-invasive
Easy-to-use**
↑



REFERENCES

1) American Cancer Society. Cancer Facts & Figures 2022. Atlanta: American Cancer Society; 2022. 2) Shaitelman SF, et al. Recent Progress in the Treatment and Prevention of Cancer-Related Lymphedema. CA Cancer J Clin. 2015;00:00-00. 3) Brown JC, et al. The Prevalence of Lymphedema Symptoms among Long-Term Cancer Survivors with or at-risk for Lower Limb Lymphedema. Am J Phys Med Rehabil. 2013 March; 92(3): 223-231. doi:10.1097/PHM.0b013e31826edd97. 4) Herd-Smith, A., et al., Prognostic factors for lymphedema after primary treatment of breast carcinoma. Cancer, 2001. 92(7): p. 1783-7. 5) Ridner SH, et al. A Comparison of Bioimpedance Spectroscopy or Tape Measure Triggered Compression Intervention in Chronic Breast Cancer Lymphedema Prevention. Lymphatic Research and Biology 2022. <https://doi.org/10.1245/s10434-019-07344-5>.