


Improving Survivorship for Breast Cancer Patients

LATEST RESEARCH SHOWS LYMPHEDEMA IS A RISK OF BREAST CANCER TREATMENT



newly diagnosed breast cancer patients every year in the U.S.¹



of breast cancer patients are at risk of developing arm lymphedema²



breast cancer survivors will be impacted by arm lymphedema³

Early Detection & Intervention are Key to Preventing Lymphedema



of patients did not progress to chronic lymphedema with early detection using L-Dex & intervention through 3 years⁴

The opportunity to prevent lymphedema is before noticeable swelling exists

Testing for Early Lymphedema is Easy

L-Dex® testing with the SOZO® Digital Health Platform



Learn more >>> www.impedimed.com/preventing-lymphedema



REFERENCES

- 1) American Cancer Society. Cancer Facts & Figures 2022. Atlanta: American Cancer Society; 2022.
- 2) American Cancer Society. Cancer Treatment & Survivorship Facts & Figures 2019–2021. Atlanta: American Cancer Society; 2019.
- 3) Gillespie TC, et al. Breast cancer-related lymphedema: risk factors, precautionary measures, and treatments. *Gland Surg* 2018;7(4):379-403.
- 4) Ridner SH, et al. A Comparison of Bioimpedance Spectroscopy or Tape Measure Triggered Compression Intervention in Chronic Breast Cancer Lymphedema Prevention. *Lymphatic Research and Biology* 2022.