

Improving Survivorship for Breast Cancer Patients



310

newly diagnosed breast cancer patients every year in the U.S.¹



82%

of breast cancer patients are at risk of developing arm lymphedema²



1 in 5

breast cancer survivors will be impacted by arm lymphedema³

Early Detection & Intervention are Key to Preventing Lymphedema

92% ↓

of patients did not progress to chronic lymphedema with early detection using L-Dex and intervention through 3 years⁴

The opportunity to prevent lymphedema is before noticeable swelling exists

Testing for Early Lymphedema is Easy

L-Dex® testing with the SOZO® Digital Health Platform



REFERENCES

- 1) Breast Cancer.org 2024 American Cancer Society, Inc.
- 2) American Cancer Society. Cancer Treatment & Survivorship Facts & Figures 2019-2021. Atlanta: American Cancer Society; 2019.
- 3) Gillespie TC, et al. Breast cancer-related lymphedema: risk factors, precautionary measures, and treatments. Gland Surg 2018;7(4):379-403.
- 4) Ridner SH, et al. A Comparison of Bioimpedance Spectroscopy or Tape Measure Triggered Compression Intervention in Chronic Breast Cancer Lymphedema Prevention. Lymphatic Research and Biology 2022.

LEARN MORE

